



PEROXigard™

One-Step Disinfectant Cleaner & Deodorizer
Ready to Use

FOR USE IN LIFE SCIENCES

For Use in Life Science Research Facilities, Vivariums, Laboratories, and Animal Research Facilities.

For disinfection and cleaning of hard non-porous surfaces found in animal premises facilities such as biosafety cabinets, lab benches, animal transportation vehicles, animal housing facilities, animal examination rooms, animal imaging and diagnostics rooms, quarantine facilities, and general laboratory equipment.

Claims

A Powerful Disinfectant Effective Against:

TUBERCULOCIDAL: 3 MIN.

Mycobacterium terrae

VIRUCIDAL: 3 MIN.

Proven effectiveness against the Poliovirus Type 1, Sabin strain type 1 which allows for a Broad-Spectrum Virucide claim against most enveloped and non-enveloped viruses.

Adenovirus type 5

Canine Parvovirus (CPV), the Cornell strain

Feline Calicivirus, F9 Strain, as a surrogate for Norwalk and Norwalk-like viruses

HIV-1 Human Immunodeficiency Virus (HIV), Strain HTLV-IIIB (HIV-1)

Human Coronavirus 229E

Human Parainfluenza Virus 3

Influenza Virus PR8 strain

Murine Norovirus type 1 (MNV-1) strain S99)

Rhinovirus serotype 14

Rotavirus Wa strain

This product has demonstrated effectiveness against Poliovirus and is expected to inactivate all Influenza A viruses including 2009 (H1N1) pandemic Influenza A virus

BACTERICIDAL: 3 MIN.

Enterobacter aerogenes

Escherichia coli O157:H7

Methicillin-resistant Staphylococcus aureus (MRSA)

Pseudomonas aeruginosa

Salmonella enterica

Staphylococcus aureus

Vancomycin-resistant Enterococcus faecalis (VRE)



Reference Sheet

Manufactured by:
Virox® Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578 | virox.com



READ SAFETY DATA SHEET BEFORE USING PRODUCT
SDS available online at www.virox.com

FUNGICIDAL: 3 MIN.

Candida albicans

Trichophyton mentagrophytes

BROAD-SPECTRUM SANITIZING: 30 SEC.

Acinetobacter baumannii

Campylobacter jejuni

Escherichia coli O157:H7

Klebsiella pneumoniae

Methicillin-resistant Staphylococcus aureus (MRSA)

Pseudomonas aeruginosa

Salmonella enterica

Staphylococcus aureus

Vancomycin-resistant Enterococcus faecalis (VRE)

Germicidal activity of this product was determined in accordance with the Canadian General Standards Board's standard CAN/CGSB-2.161-97

Disinfectant Use Directions

HEAVILY SOILED SURFACES REQUIRE CLEANING PRIOR TO DISINFECTION

Cleaning, Disinfection and Deodorizing of Laboratory Equipment and Hard Non-Porous Surfaces: Pre-clean heavily soiled areas. Apply solution to surface until thoroughly wet. Ensure surface remains wet for 3 minutes. Allow to air dry.

Directions for Use in Animal Containment Areas

1. Remove all animals and their feed from premises, and enclosures prior to disinfection.
2. Remove heavy soil such as urine and manure from the surfaces of facilities and fixtures occupied or traversed by animals.
3. Empty or remove all feeding and watering equipment.
4. Thoroughly clean all surfaces with detergent or product and rinse with potable water.
5. Apply product to all surfaces for a period of 3 minutes.
6. All equipment used in the handling and restraining of animals, and for removing litter and manure, should be immersed in solution for 3 minutes. Allow items to dry before re-using.
7. Ventilate closed spaces. Do not house animals or employ equipment until the product has been absorbed, set, or dried.
8. Thoroughly scrub all treated surfaces or equipment that come in contact with feed and drinking water such as automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

Directions for Use in Food Processing Establishments

Apply solution to surface with cloth or disposable wipe. Ensure surface remains wet for 3 minutes. For surfaces and/or equipment that may come in contact with food, a rinse with potable water is required.

Broad-Spectrum Sanitizing on Environmental Surfaces (>99.9%):

Apply to surface, allow to remain wet for 30 seconds. Allow to air dry. Food contact surfaces require rinsing with potable water.

Special Instructions for Cleaning and Decontamination

Against HIV (Human Immunodeficiency Virus) on objects and surfaces soiled with blood/body fluids. This product is intended for use against HIV only in those settings where the virus would be expected to be encountered, such as settings where contamination by blood or body fluids is likely.

Personal Protection: Disposable gloves, gowns, face masks, or eye covering as appropriate, must be worn during all cleaning of body fluids, blood and decontamination procedures.

Cleaning and Disinfecting Surfaces of Blood and Body

Fluids: Gloves should be worn. Remove excess blood and fluid with absorbent materials. Clean contaminated area: Apply to surface, allow surface to remain wet for 30 seconds, wipe dry. Disinfect contaminated area: Apply to surface, allow surface to remain wet for 3 minutes.

Disposal of Infectious Material: Products contaminated with blood or body fluids should be disposed of according to Federal, Provincial, and local regulations for infectious waste disposal.

Precautionary Statements

No Personal Protective Equipment is required when handling this product. **KEEP OUT OF REACH OF CHILDREN.** Avoid storage at elevated temperatures. Avoid contamination of food in the application and storage of this product. Do not mix with other cleaning or disinfecting products.

FIRST AID: If in contact with eyes, flush immediately and thoroughly with water. Call a physician if irritation persists. If ingested, do not induce vomiting. Call a physician or a poison control center immediately. **TAKE CONTAINER, LABEL OR PRODUCT NAME AND DIN WITH YOU WHEN SEEKING MEDICAL ATTENTION.**

STORAGE: Avoid storage at elevated temperatures. Store at room temperature.

DISPOSAL: For information on the disposal of unused, unwanted product and the clean up of spills, contact the Provincial Regulatory Agency or the Manufacturer. Rinse the container with water. Dispose of container in accordance with municipal, provincial and federal regulations. Do not reuse containers.